

THE $M|M|\infty$ QUEUE TRANSIENT PROBABILITIES

Manuel Alberto M. Ferreira

Instituto Universitário de Lisboa (ISCTE-IUL), Business Research Unit (BRU-IUL) and Information Sciences, Technologies and Architecture Research Center (ISTAR-IUL), PORTUGAL

E-mail: manuel.ferreira@iscte.pt

ABSTRACT

In the theory and applications of queuing systems, transient probabilities play a key role. Often, due to the difficulty of their calculation they are replaced by the stationary probabilities. In the case of $M|G|\infty$ systems, those calculations are much friendlier than usual. We present in this work formulas for the transient probabilities of the $M|M|\infty$ system, where are considered service times with exponential distribution, showing that it is also possible to calculate them with initial conditions different from those usually considered.

Keywords: Transient probabilities, $M|M|\infty$, busy period, exponential distribution.

Mathematics Subject Classification: 60G99

INTRODUCTION

In the $M|G|\infty$ queue, customers arrive according to a Poisson process at rate λ , upon its arrival receive immediately a service with time length *d. f.* $G(\cdot)$ and mean α . The traffic intensity is $\rho = \lambda\alpha$, see for instance [7].

Call $N(t)$ the number of occupied servers, at time $t, t \geq 0$ in the $M|G|\infty$ queue. Define the transient probabilities $p_{0n}(t)$ as

$$p_{0n}(t) = P\{N(t) = n | N(0) = 0\}, n = 0, 1, 2, \dots, t \geq 0 \quad (1.1)$$

So, see [7],

$$p_{0n}(t) = \frac{\left(\lambda \int_0^t [1 - G(v)] dv\right)^n}{n!} e^{-\lambda \int_0^t [1 - G(v)] dv}, n = 0, 1, 2, \dots$$

The stationary probabilities are the limit probabilities and so

$$p_{0n} = \lim_{t \rightarrow \infty} p_{0n}(t) = \frac{\rho^n}{n!} e^{-\rho}, n = 0, 1, 2, \dots \quad (1.2)$$

since $\alpha = \int_0^\infty [1 - G(v)] dv$ because $G(\cdot)$ is a positive distribution *d. f.*

As it happens for any queue, in the $M|G|\infty$ queue activity there is a sequence of idle and busy periods. For this queue the study of the busy period length distribution is very important since, as it is part of its definition, a customer must find immediately an available server upon its arrival. So, it is important for the manager to know how many, and how long, servers must be in prevention, see [1] and [5, 6].

Define now

$$p'_{1'n}(t) = P\{N(t) = n | N(0) = 1'\}, n = 0, 1, 2, \dots, t \geq 0 \quad (1.3)$$

meaning $N(0) = 1'$ that the time origin is considered in an instant at which a customer arrives at the $M|G|\infty$ system finding it empty, that is: in the beginning of a busy period. At the instant $t \geq 0$ it may occur:

- Either the customer that arrived at the initial instant abandoned the system, with probability $G(t)$, or goes on being served, with probability $1 - G(t)$,
- The other servers, that were unoccupied at the time origin, are either unoccupied or occupied with $1, 2, \dots$ customers, with probability $p_{0n}(t), n = 0, 1, 2, \dots$ given by (1.1).

As both systems, the one of the initial customer and the other of the initially unoccupied servers, are independent, consequently:

$$\begin{aligned} p'_{1'0}(t) &= p_{00}(t)G(t) \\ p'_{1'n}(t) &= p_{0n}(t)G(t) + p_{0n-1}(t)(1 - G(t)), n = 1, 2, \dots \end{aligned} \quad (1.4).$$

The stationary probabilities are also

$$p'_{1'n} = \lim_{t \rightarrow \infty} p'_{1'n}(t) = \frac{\rho^n}{n!} e^{-\rho}, n = 0, 1, 2, \dots \quad (1.5).$$

THE $M|M|\infty$ CASE

For the $M|M|\infty$ queue, that is exponential service times, (1.1) becomes:

$$p_{0n}^M(t) = \frac{\left(\rho \left(1 - e^{-\frac{t}{\alpha}}\right)\right)^n}{n!} e^{-\rho \left(1 - e^{-\frac{t}{\alpha}}\right)}, n = 0, 1, 2, \dots \quad (2.1)$$

and (1.4)

$$\begin{aligned} p_{10}^M(t) &= p_{00}^M(t) \left(1 - e^{-\frac{t}{\alpha}}\right) \\ p_{1n}^M(t) &= p_{0n}^M(t) \left(1 - e^{-\frac{t}{\alpha}}\right) + p_{0n-1}^M(t) e^{-\frac{t}{\alpha}}, n = 1, 2, \dots \end{aligned} \quad (2.2),$$

owing to the exponential distribution lack of memory, that allows to consider as initial instant anyone at which there is only one customer in the system, not necessarily demanding it is the beginning of a busy period.

Analogously to the procedure followed to deduce (1.4), considering two independent subsystems, one with the clients present at the initial instant and another with the infinite servers vacated at that instant, it is concluded, taking into account the memory shortage of the exponential distribution that for the $M|M|\infty$ queue:

$$\begin{aligned} p_{mn}^M(t) &= \sum_{k=0}^m \binom{m}{k} \frac{\left(\rho \left(1 - e^{-\frac{t}{\alpha}}\right)\right)^{n-k}}{(n-k)!} e^{-kt/\alpha - \rho \left(1 - e^{-\frac{t}{\alpha}}\right)} \left(1 - e^{-\frac{t}{\alpha}}\right)^{m-k} I \\ &(n-k), m = 0, 1, 2, \dots, n = 0, 1, 2, \dots \text{ and } I(n-k) = \begin{cases} 1, n \geq k \\ 0, n < k \end{cases} \end{aligned} \quad (2.3).$$

In particular, for $n = 0$:

$$p_{m0}^M = e^{-\rho\left(1-e^{-\frac{t}{\alpha}}\right)} \left(1 - e^{-\frac{t}{\alpha}}\right)^m, m = 0,1,2, \dots (2.4).$$

Theorem 2.1

For the $M|M|\infty$ queue:

i) $m \geq \rho$

$p_{m0}^M(t), t > 0$ is an increasing function,

ii) $0 < m < \rho$

a) $p_{m0}^M(t)$, is an increasing function in $]0, t_m[$

b) $p_{m0}^M(t)$, is a decreasing function in $]t_m, 0[$

c) The $p_{m0}^M(t)$ maximum is

$$p_{m0}^M(t_m) = \left(\frac{m}{\rho e}\right)^m (2.5)$$

being

$$t_m = -\alpha \ln\left(1 - \frac{m}{\rho}\right) (2.6),$$

iii) $m=0$

$p_{m0}^M(t), t > 0$ is a decreasing function.

Dem: It is enough to have in mind that $\frac{d}{dt}$

$$p_{m0}^M(t) = e^{-\rho\left(1-e^{-\frac{t}{\alpha}}\right)} \left(1 - e^{-\frac{t}{\alpha}}\right)^m e^{-\frac{t}{\alpha}} \left(\frac{m}{\alpha} \frac{1}{1-e^{-\frac{t}{\alpha}}} - \lambda\right).$$

Obs: If $\rho < 1$, $p_{m0}^M(t), m = 1,2, \dots$ always can be normalized in order to behave as d.f. ■

CONCLUSIONS

In this text, the transient probabilities for the $M|G|\infty$ queue are reviewed, and clarified for the $M|M|\infty$ queue. Based on this are determined transient probabilities for the $M|M|\infty$ system, with initial conditions different from the usual, where play a key role the exponential distribution lack of memory.

REFERENCES

- 1) J.A. Filipe, M.A.M. Ferreira, Infinite Servers Queue Systems Busy Period-A Practical Case in Logistics Problems Solving. Applied Mathematical Sciences, 9(2015), 25-28, 1221-1228.
- 2) J. Walrand, An Introduction to Queueing Networks. New Jersey: Prentice-Hall, Inc. 1988.
- 3) L. Tackác, An Introduction to Queueing Theory. Oxford University Press. New York, 1962.
- 4) M.A.M. Ferreira, Momentos da Duração do Período de Ocupação de Sistemas de Filas de Espera com Infinitos servidores. Actas da 5ª Conferência CEMAPRE, 429-438. ISEG, Lisboa 26, 27, 28 de Maio, 1997.

- 5) M.A.M. Ferreira, The Exponentiality of the $M|G|\infty$ queue busy period. XII Jornadas Luso-Espanholas de Gestão Científica-ACTAS (Volume VIII-Economia da Empresa e Matemática Aplicada), 267-272. Universidade da Beira Interior, Departamento de Gestão e Economia, Covilhã, Abril 2002.
- 6) M.A.M. Ferreira, J. A. Filipe, Occupation Study of a Surgery Service through a Queues Model. Applied Mathematical Sciences, 9(2015), 93-96, 4791-4798.
- 7) M.A.M. Ferreira, M. Andrade, The Ties between the $M|G|\infty$ Queue System Transient Behavior and the Busy Period. International Journal of Academic Research, 1 (2009),1, 84-92.
- 8) M.A.M. Ferreira, M. Andrade, Looking to a $M|G|\infty$ system occupation through a Riccati equation. Journal of Mathematics and Technology, 1(2010), 2, 58-62.
- 9) M.A.M. Ferreira, M. Andrade, $M|G|\infty$ System Transient Behavior with Time Origin at the Beginning of a Busy Period Mean and Variance. APLIMAT- Journal of Applied Mathematics, 3(2010), 3, 213-221.
- 10) M.A.M. Ferreira, M. Andrade, $M|G|\infty$ Queue Busy Period Tail. Journal of Mathematics and Technology, 1(2010), 3, 11-16.
- 11) M.A.M. Ferreira, M. Andrade, Algorithm for the Calculation of the
- 12) Laplace-Stieltjes Transform of the Sojourn Time of a Costumer in an Open Network of Queues, with a Product Form Equilibrium Distribution, assuming Independent Sojourn Times in each Node. Journal of Mathematics and Technology, 1(2010), 4, 31-36.
- 13) M.A.M. Ferreira, M. Andrade, Infinite Servers Queue Systems Busy Period Length Moments. Journal of Mathematics and Technology, 3(2012), 2, 4-7.
- 14) M.A.M. Ferreira, M. Andrade, J. A Filipe, The Riccati Equation in the $M|G|\infty$ System Busy Cycle Study. Journal of Mathematics, Statistics and Allied Fields, 2 (2008), 1.
- 15) M.A.M. Ferreira, The Laplace transform in the context of some of the queue related probability distributions. Acta Scientiae et Intellectus, 4(2018), 4, 24-31.
- 16) M.J. Carrillo. Extensions of Palm's Theorem: A Review. Management Science, 37(1991), 6, 739-744.
- 17) R.L. Disney and D. König. Queueing networks: a survey of their random processes. Siam Review, 3(1985), 335-403.
- 18) S. Skiena, "Polya's Theory of Counting." §1.2.6 in Implementing Discrete Mathematics: Combinatorics and Graph Theory with Mathematica. Reading, MA: Addison-Wesley, pp. 25-26, 1990.
- 19) W. Stadje. The Busy Period of the Queueing System $M|G|\infty$. Journal of Applied Probability, 22(1985), 697-704.